arXiv:math/0412058v1 [math.DS] 2 Dec 2004

On holomorphic foliations with projective transverse structure

Bruno Scárdua

Abstract

We study codimension one holomorphic foliations on complex projective spaces and compact manifolds under the assumption that the foliation has a projective transverse structure in the complement of some invariant codimension one analytic subset. The basic motivation is the characterization of pull-backs of Riccati foliations on projective spaces. Our techniques apply to give a description of the generic models of codimension one foliations on compact manifolds of dimension ≥ 3.

1 Introduction and summary

Let $\mathcal{F}$ be a codimension one holomorphic foliation on a connected complex manifold M^m, of dimension $m \geq 2$, having singular set $\operatorname{sing}(\mathcal{F})$ of codimension ≥ 2. The foliation $\mathcal{F}$ is *transversely projective* if there is an open cover $\{U_j,\, j \in J\}$ of $M \setminus \operatorname{sing}(\mathcal{F})$ for which in each U_j the foliation is given by a holomorphic submersion $f_j \colon U_j \to \mathbb{C}$ such that in each intersection $U_i \cap U_j \neq \emptyset$ we have $f_i = \frac{a_{ij} f_j + b_{ij}}{c_{ij} f_j + d_{ij}}$ for some locally constant functions $a_{ij}, b_{ij}, c_{ij}, d_{ij}$ with $a_{ij}d_{ij} - b_{ij}c_{ij} = 1$. If we have $f_i = a_{ij} f_j + b_{ij}$ for locally constant functions $a_{ij} \neq 0$, b_{ij} then $\mathcal{F}$ is *transversely affine* in M ([10]). The basic references for transversely affine and transversely projective foliations are found in [5]. In general a holomorphic foliation on a rational projective manifold cannot be transversely projective on the whole manifold, nevertheless there are many interesting examples where the foliation admits a projective transverse structure in the complement of an invariant algebraic subset of codimension one. Samples of this situation are given by logarithmic foliations and Bernoulli foliations (for the transversely affine case) and Riccati foliations (for the transversely projective case) on the complex projective plane $\mathbb{C}P(2)$. Such foliations are considered in [10], [11] and [3] where classification results are found. These results regard mainly the transversely affine case which is proved to be strongly linked to logarithmic and Bernoulli foliations if some hypothesis on the singular set of the foliation is made. Nevertheless, in [4] it is given an example of a foliation on $\mathbb{C}P(2)$ which is transversely projective in the complement of an algebraic invariant curve and which is neither logarithmic nor a Bernoulli or a Riccati pull-back. One of the main results in [4] states, in simple words, that *if M is compact of dimension $m \geq 3$ and $\mathcal{F}$ admits a meromorphic vector field X in M which is generically transverse to $\mathcal{F}$ then we have two possibilities:* (1) *$\mathcal{F}$ is the meromorphic pull-back of an algebraic foliation in a projective manifold.* (2) *$\mathcal{F}$ is transversely projective in $M \setminus \Lambda$ for some compact analytic invariant subset $\Lambda \subset M$ of codimension one.* The above result gives a strong property for foliations on compact manifolds of dimension $m \geq 3$ and motivates the study of transversely projective foliations using transcendental techniques. This is one of the motivations for this work. One second motivation is to resume the classification of transversely projective foliations on projective spaces studied in [10]. Our approach is based in (trancendental) techniques of extension of the transverse structure through the singularities of the foliation lying in the subset $\Lambda \subset M$. This sort of idea was already present in [10],[3] but we must now extend a different object; the transverse foliation associated to a projective transverse structure. The rough idea is to use the existence of separatrices transverse to Λ in order to begin the extension procedure. As a sample of our results we have:

Theorem 1. *Let $\mathcal{F}$ be a codimension one holomorphic foliation on $\overline{\mathbb{C}} \times \overline{\mathbb{C}}$. Assume that $\mathcal{F}$ is transversely projective in the complement of an invariant algebraic curve $\Lambda \subset \overline{\mathbb{C}} \times \overline{\mathbb{C}}$ and that the singularities of $\mathcal{F}$ in $\overline{\mathbb{C}} \times \overline{\mathbb{C}}$ are hyperbolic. Then $\mathcal{F}$ is a logarithmic foliation or a rational pull-back of a Riccati foliation.*

We point-out that (cf. Example 1): (i) A generic Riccati foliation on $\overline{\mathbb{C}} \times \overline{\mathbb{C}}$ has only hyperbolic singularities and satisfies the above hypotheses. (ii) We have a similar statement for foliations on $\mathbb{C}P(2)$ which are transversely projective in the complement of an algebraic invariant curve. Nevertheless, we have to take into account the fact that a Riccati foliation pull-back on $\mathbb{C}P(2)$ exhibits some nonhyperbolic singularities (see Example 1 for a detailed description of this case). (iii) Theorem 1 admits a much sharper statement in terms of the Reduction of Singularities for $\mathcal{F}$ in $\operatorname{sing}(\mathcal{F}) \cap \Lambda$ (see Definition 3 and Theorems 3 and 4).

From the Algebraic Reduction Theorem in [4] (see Theorem 5) we obtain:

Theorem 2. *Let $\mathcal{F}$ be a pseudo-parallelizable holomorphic foliation on a simply-connected compact manifold M of dimension ≥ 3. Assume that the singularities of $\mathcal{F}$ in M are of types:* (1) *singularities belonging to codimension ≥ 3 components of* $\operatorname{sing}(\mathcal{F})$ *and* (2) *hyperbolic singularities. Then we have the following possibilities:*

(i) *$\mathcal{F}$ is pseudo-algebraic: there is a meromorphic map $\tau \colon M \to N$, where N is an algebraic projective manifold, such that $\mathcal{F}$ is the pull-back by τ of an algebraic foliation on N.*

(ii) *$\mathcal{F}$ admits an extended affine transverse structure in M in the sense of* [10]: *$\mathcal{F}$ is given by a meromorphic one-form Ω on M for which there is a closed meromorphic one-form η with simple poles on M such that $d\Omega = \eta \wedge \Omega$.*

(iii) *$\mathcal{F}$ is a meromorphic pull-back of a Riccati foliation on $\overline{\mathbb{C}} \times \overline{\mathbb{C}}$.*

(iv) *$\mathcal{F}$ is a logarithmic foliation, i.e., $\mathcal{F}$ is given by a closed meromorphic one-form with simple poles in M.*

In case (ii) $\mathcal{F}$ is transversely affine in $M \backslash \Lambda$ where $\Lambda \subset M$ is the compact codimension one analytic subset $\Lambda = (\eta)_\infty$. Since η is meromorphic in M with simple poles $\mathcal{F}$ admits an *extended affine transverse structure* in M according to the terminology in [10]. In particular, we know exactly how to describe the holonomy group of the leaves in Λ in terms of the residues of η in Λ (see [10] Lemma 3.1 § 3). We have a Liouvillian first integral F for $\mathcal{F}$ given by $F = \int \frac{\Omega}{e^{\int \eta}}$ and we believe that more can be said in this case (see the main result of [3] for instance). We can relax the hypotheses on the singularities of $\mathcal{F}$ in Theorem 2 by allowing singularities that have a local integrating factor of holomorphic type but which do not exhibit meromorphic first integrals.

Theorem 3. *Let $\mathcal{F}$ be a holomorphic foliation on $\mathbb{C}P(2)$ and transversely projective outside an algebraic invariant curve $\Lambda \subset \mathbb{C}P(2)$. Assume that $\mathcal{F}$ has noncentral separatrices transverse to Λ* (*in the sense of* Definition 3)*. Then $\mathcal{F}$ is a rational pull-back of a Riccati foliation.*

2 Projective transverse structures and projective triples

According to [10] a holomorphic foliation $\mathcal{F}$ of codimension one and having singular set $\operatorname{sing}(\mathcal{F})$ of codimension ≥ 2 in a complex manifold M^m, $m \geq 2$ is *transversely projective* if the underlying nonsingular foliation $\mathcal{F}' = \mathcal{F}|_{M \backslash \operatorname{sing}(\mathcal{F})}$ is transversely projective in $M' = M \backslash \operatorname{sing}(\mathcal{F})$.

Example 1 (Riccati Foliations). Fix affine coordinates $(x,y) \in \mathbb{C}^2$ and consider a polynomial vector field $X(x,y) = p(x)\,\frac{\partial}{\partial x} + \big(a(x)y^2 + b(x)y + c(x)\big)\,\frac{\partial}{\partial y}$ on $\mathbb{C}^2$. Then X defines a *Riccati foliation* $\mathcal{F}_X$ on $\overline{\mathbb{C}} \times \overline{\mathbb{C}}$ as follows: if we change coordinates via $u = \frac{1}{x}$, $\ v = \frac{1}{y}$ then we obtain $X(x,v) = p(x)\,\frac{\partial}{\partial x} - \big(a(x) + b(x)v + c(x)v^2\big)\,\frac{\partial}{\partial v}\cdot$ Similarly for $X(u,y) = u^{-n}[\tilde{p}(u)\,\frac{\partial}{\partial u} + \big(\tilde{a}(u)y^2 + \tilde{b}(u)y + \tilde{c}(u)\big)\,\frac{\partial}{\partial y}]$ and $X(u,v) = u^{-n}[\tilde{p}(u)\,\frac{\partial}{\partial u} + \big(\tilde{a}(u) + \tilde{b}(u)v + \tilde{c}(u)v^2\big)\frac{\partial}{\partial v}]$. The similarity of these four expressions shows that X defines a holomorphic foliation $\mathcal{F}_X$ with isolated singularities on $\overline{\mathbb{C}} \times \overline{\mathbb{C}}$ and having a geometry as follows (see Figure 1):
(i) $\mathcal{F}_X$ is transverse to the fibers $\{a\} \times \overline{\mathbb{C}}$ except for invariant fibers which are given in $\mathbb{C}^2$ by $\{p(x) = 0\}$.
(ii) If $\Lambda = \bigcup\limits_{j=1}^{r} \{a_j\} \times \overline{\mathbb{C}}$ is the set of invariant fibers then $\mathcal{F}_X$ is transversely projective in $(\overline{\mathbb{C}} \times \overline{\mathbb{C}})\backslash\Lambda$. Indeed, $\mathcal{F}_X|_{(\overline{\mathbb{C}}\times\overline{\mathbb{C}})\backslash\Lambda}$ is conjugate to the suspension of a representation $\varphi\colon \pi_1(\overline{\mathbb{C}}\backslash \bigcup\limits_{j=1}^{r}\{a_j\}) \to \mathbb{P}\,\mathrm{SL}(2,\mathbb{C})$.
(iii) For a generic choice of the coefficients $a(x), b(x), c(x), p(x) \in \mathbb{C}[x]$ the singularities of $\mathcal{F}_X$ on $\overline{\mathbb{C}} \times \overline{\mathbb{C}}$ are hyperbolic, Λ is the only algebraic invariant set and therefore for each singularity $q \in \operatorname{sing}(\mathcal{F}_X) \subset \Lambda$ there is a local separatrix of $\mathcal{F}_X$ transverse to Λ passing through q.

Now we consider the canonical way of passing from $\overline{\mathbb{C}}\times\overline{\mathbb{C}}$ to $\mathbb{C}P(2)$ by a map $\sigma\colon \overline{\mathbb{C}}\times\overline{\mathbb{C}} \to \mathbb{C}P(2)$ obtained as a sequence of birrational maps as follows: first blow-up a point, for example the origin, of $\mathbb{C}^2 \subset \overline{\mathbb{C}} \times \overline{\mathbb{C}}$ then blow-down two suitable projective lines of self-intersection equals -1 as indicated in Figure 1. Following this process step by step we conclude that the foliation $\mathcal{F} = \sigma_*(\mathcal{F}_X) = (\sigma^{-1})^*(\mathcal{F}_X)$ induced by $\mathcal{F}_X$ on $\mathbb{C}P(2)$ has the following characteristics:
(i') $\mathcal{F}$ is transversely projective in $\mathbb{C}P(2)\backslash\Lambda$ where $\Lambda \subset \mathbb{C}P(2)$ is the union of a finite number of projective lines of the form $\bigcup\limits_{j=1}^{r} \overline{\{x = a_j\}} \subset \mathbb{C}P(2)$ in a suitable affine chart $(x,y) \in \mathbb{C}^2 \subset \mathbb{C}P(2)$.
(ii') For a generic choice of the coefficients of X, the singularities of $\mathcal{F}$ in Λ are hyperbolic except for one single dicritical singularity $q_\infty\colon (x = \infty, y = 0) \in \mathbb{C}P(2)$ which after one blow-up gives a nonsingular foliation transverse to the projective line except for a single tangency point. The foliation $\mathcal{F}$ also has two other nonhyperbolic singularities, belonging to the line at infinity $L_\infty = \mathbb{C}P(2) \setminus \mathbb{C}^2$, which is invariant, one linearizable with holomorphic first integral and the other dicritical of radial type, admitting a meromorphic first integral. Also, in general, $\Lambda \cup L_\infty$ is the only algebraic invariant set and $\operatorname{sing}(\mathcal{F}) \subset \Lambda \cup L_\infty$.
(iii') Finally, we stress that on $\mathbb{C}P(2)$ the foliation $\mathcal{F}$ is transversely projective in a neighborhood of $L_\infty \setminus (L_\infty \cap \operatorname{sing}(\mathcal{F}))$.

Now we give a formulation of the existence of a projective transverse structure for a given foliation in terms of differential forms. Suppose that $\operatorname{sing}(\mathcal{F}) = \emptyset$ and $m = 2$ and let $\mathcal{F}$ be given by a holomorphic one-form Ω with $\Omega \neq 0$. Also assume there is a holomorphic one-form η such that $d\Omega = \eta \wedge \Omega$.

Proposition 1 ([10]). *Under the above hypotheses the following conditions are equivalent:* (i) *$\mathcal{F}$ is transversely projective.* (ii) *There exists a holomorphic one-form ξ such that $d\eta = \Omega \wedge \xi$ and $d\xi = \xi \wedge \eta$.*

Above we deal with the holomorphic nonsingular case nevertheless we also have:

Lemma 1 ([10]). *The triple (Ω, η, ξ) defining a projective transverse structure for $\mathcal{F}$ can be modified preserving this property as: $\Omega' = g\Omega$; $\ \eta' = \eta + \frac{dg}{g} + h\Omega$; $\ \xi' = \frac{1}{g}(\xi - dh - h\eta - \frac{h^2}{2}\,\Omega)$ where g and h are holomorphic functions.*



Figure 1: A Riccati foliation from $\overline{\mathbb{C}} \times \overline{\mathbb{C}}$ to $\mathbb{C}P(2)$.

This allows to consider the meromorphic and singular case for Ω. In particular we can modify (Ω, η, ξ) into $(g\Omega,\ \eta + \frac{dg}{g}\ ,\ \frac{1}{g}\,\xi)$ or also $(\frac{1}{g}\,\Omega,\ \eta - \frac{dg}{g}\ ,\ g\xi)$. We obtain therefore the following:

Definition 1 ([3]). Let $\mathcal{F}$ be a foliation given by a meromorphic one-form Ω on M with isolated singularities and such that $(\Omega)_\infty$ has no invariant component. We say that Ω, η, ξ is a order one *projective triple* if η, ξ are meromorphic one-forms on M such that:
(i) $d\Omega = \eta \wedge \Omega$, $(\eta)_\infty = (\Omega)_\infty$, $(\eta)_\infty$ has order one and for any irreducible component $C \subset (\Omega)_\infty$ we have that $\eta + k\frac{df}{f}$ is holomorphic where $\{f = 0\}$ in a reduced local equation of C and $k = \operatorname{ord}\big((\Omega)_\infty, C\big)$ is the order of $(\Omega)_\infty$ along C.
(ii) $d\eta = \Omega \wedge \xi$ and $d\xi = \xi \wedge \eta$ where ξ is holomorphic and has zero divisor $(\xi)_0 = (\Omega)_\infty$ and for each irreducible component $C \subset (\Omega)_\infty$ we have $\operatorname{ord}\big((\xi)_0, C\big) = \operatorname{ord}\big((\Omega)_\infty, C\big)$.

In general we also allow triples which come from a order one projective triple Ω, η, ξ by modifications of the form $\Omega' = g\Omega$, $\eta' = \eta + \frac{dg}{g} + h\Omega$, $\xi' = \frac{1}{g}\xi - dh - h\eta - \frac{h^2}{2} \cdot$ Such a triple will be called a *meromorphic projective triple* and still defines a projective transverse structure for $\mathcal{F}$. Notice also that given a projective triple (Ω, η, ξ) the one-form ξ also defines a foliation with a projective transverse structure given by the projective triple $(\xi, -\eta, \Omega)$; we will usually denote this *transverse foliation* by $\mathcal{F}^\perp$. For the case of a Riccati foliation (see Example 1 above) we can for instance take $\Omega = p(x)dy - (y^2\,c(x) - yb(x) - a(x))dx$, $\eta = 2\frac{dy}{y} + \frac{p'+b}{p}\,dx + \frac{2a}{yp}\,dx$ and $\xi = \frac{-2a}{y^2p^2}\,dx$ and modify it into $\Omega = pdy - (y^2a + by + c)dx$, $\eta = \frac{p'+b+2ay}{p}\,dx$, $\xi = \frac{2a}{p^2}\,dx$. We shall also use the following lemma. Let (Ω, η, ξ) be meromorphic one-forms which satisfy the *projective relations*, i.e., such that $d\Omega = \eta \wedge \Omega$, $\; d\eta = \Omega \wedge \xi$ and $d\xi = \xi \wedge \eta$.

Lemma 2 ([10]). *Given a meromorphic one-form ξ' the triple (Ω, η, ξ') also satisfies the projective relations, i.e., we have $d\eta = \Omega \wedge \xi'$ and $d\xi' = \xi' \wedge \eta$ if, and only if $\xi' = \xi + F\Omega$ for some meromorphic function F such that $d\Omega = -\frac{dF}{2F} \wedge \Omega$.*

We resume the study of the Lie Algebra associated to a Riccati foliation as in Example 1. Let $\Omega := p(x)dy - \big(y^2 a(x) + b(x)y + c(x)\big)dx$ be the natural polynomial one-form defining $\mathcal{F}$ in $\mathbb{C}^2$ with affine coordinates (x, y). Put $\eta := \frac{p'(x)+b(x)+2ya(x)}{p(x)}\,dx$ then η is rational with polar set such that $(\eta)_\infty \cap \mathbb{C}^2 = \Lambda \cap \mathbb{C}^2$. We have $d\Omega = \eta \wedge \Omega$. Finally, define $\xi := \frac{2a(x)}{p(x)^2}\,dx$. Then we have $d\eta = \Omega \wedge \xi$ and $d\xi = \xi \wedge \eta$. Also ξ defines the rational fibration $\overline{x}\colon \mathbb{C}P(2) \to \overline{\mathbb{C}}$ obtained by extending the coordinate projection $x\colon \mathbb{C}^2 \to \mathbb{C}$ which exhibits Λ as union of fibers. The forms Ω, η, ξ define a projective triple. The triple (Ω, η, ξ) is holomorphic in $\mathbb{C}^2 \backslash \Lambda$ nevertheless the projective transverse structure of the foliation $\mathcal{F}^\perp$ defined by ξ extends from $\mathbb{C}^2 \backslash \Lambda$ to all $\mathbb{C}P(2)$. Indeed, $\mathcal{F}^\perp$ coincides with the singular fibration $\overline{x}\colon \mathbb{C}P(2) \to \overline{\mathbb{C}}$ above. As we will see, this is a general fact under suitable hypothesis on the singular set of a foliation $\mathcal{F}$ on $\mathbb{C}P(2)$ admitting a projective transverse structure in the complement of an algebraic one dimensional invariant subset $\Lambda \subset \mathbb{C}P(2)$.



Figure 2: A Riccati foliation on $\mathbb{C}P(2)$ highlighting the separatrices transverse to Λ.

3 Extension of projective transverse structures

In this section we consider the following situation: $\mathcal{F}$ is a holomorphic foliation of codimension one, with singular set $\operatorname{sing}(\mathcal{F})$ of codimension two on a complex surface M^2. We have a compact analytic curve $\Lambda \subset M^2$ which is invariant by $\mathcal{F}$ and irreducible as an analytic subset of M. Denote by $\operatorname{sep}(\mathcal{F}, \Lambda)$ the set of separatrices of $\mathcal{F}$ through the singularities $q \in \operatorname{sing}(\mathcal{F}) \cap \Lambda$ and not contained in Λ. We may assume that $\operatorname{sep}(\mathcal{F}, \Lambda)$ is a codimension one (closed) analytic subset of M by replacing M by a neighborhood of Λ in M. Assume that $\operatorname{sing}(\mathcal{F}) \subset \Lambda$ and that the singularities of $\mathcal{F}$ in M are *irreducible* in the following sense (cf. § 4). Given $q \in \operatorname{sing}(\mathcal{F})$ we have a local chart $(x, y) \in U$ in a neighborhood U of q in M such that: $\Lambda \cap U : \{y = 0\}$, either $\operatorname{sep}(\mathcal{F}, \Lambda) \cap U = \emptyset$ or $\operatorname{sep}(\mathcal{F}, \Lambda) \cap U : \{x = 0\}$, $\; x(q) = y(q) = 0$ and $\mathcal{F}|_U$ is given by one of the forms below:

(Irred.i) $xdy - \lambda ydx + xy \cdot \omega_2(x, y) = 0$ for some holomorphic one-form $\omega_2(x, y)$ with $\omega_2(0, 0) = 0$ and $\lambda \in \mathbb{C} \backslash \mathbb{Q}_+$.

(Irred.ii.a) $x^{k+1}\,dy-[y(1+\lambda x^k)+A(x,y)]dx=0$ for some holomorphic function $A(x,y)$ of order $\geq k+1$, $\lambda\in\mathbb{C}$ and $k\in\mathbb{N}$. We call $\{x=0\}$ the *strong-manifold* of the singularity and shall say that the singularity *has a strong manifold transverse to* Λ.

(Irred.ii.b) $y^{k+1}\,dx-[x(1+\lambda y^k)+A(x,y)]dy=0$, where $A(x,y)$ is a holomorphic function of order $\geq k+1$, $\lambda\in\mathbb{C}$ and $k\in\mathbb{N}$. We say that $\{y=0\}$ is the *strong manifold* and that the singularity *has a central manifold transverse to* Λ.

We also assume that $\mathcal{F}$ is given by a meromorphic one-form Ω in M with isolated zeros and with polar set $(\Omega)_\infty$ in *general position* what means that $(\Omega)_\infty$ is smooth, irreducible, noninvariant by $\mathcal{F}$ and meets Λ transversely and only at points which are not singular points for $\mathcal{F}$. In particular we may assume that $(\Omega)_\infty\cap\operatorname{sep}(\mathcal{F},\Lambda)=\emptyset$. Finally, we suppose that Ω has a *logarithmic derivative adapted to* (Ω,Λ) what means the following: there exists a one-form η in M which satisfies $d\Omega=\eta\wedge\Omega$ outside $(\Omega)_\infty$, $(\eta)_\infty=(\Omega)_\infty$, $(\eta)_\infty$ has order one and if $\{f=0\}$ is a local reduced equation of $(\Omega)_\infty$ then the one-form $\eta+k\,\frac{df}{f}$ is holomorphic, where k is the order of $(\Omega)_\infty$. Write $\operatorname{sep}(\mathcal{F},\Lambda)=\bigcup\limits_{j=1}^{r}S_j$ in (disjoint) irreducible components.

Lemma 3 (Extension Lemma). *Assume that there is an irreducible component S_{j_0} of* $\operatorname{sep}(\mathcal{F},\Lambda)$ *such that $\mathcal{F}$ is transversely projective in $M\backslash(\Lambda\cup\bigcup\limits_{j\neq j_0}S_j)$. Also assume that the singularity $q_{j_0}=S_{j_0}\cap\Lambda$ is of type* (Irred.i) *or* (Irred.ii.a) (*i.e., not of type* (Irred.ii.b))*. Then $\mathcal{F}$ admits a projective triple (Ω,η,ξ) in $M\backslash\bigcup\limits_{j\neq j_0}S_j$.*

Proof: Since $\mathcal{F}$ is transversely projective in $M\backslash(\Lambda\cup\bigcup\limits_{j\neq j_0}S_j)$ according to [10] there exists a one-form ξ in $M\backslash(\Lambda\cup\bigcup\limits_{j\neq j_0}S_j)$ such that $d\eta=\Omega\wedge\xi$ and $d\xi=\xi\wedge\eta$ with (Ω,η,ξ) a projective triple on $M\backslash(\Lambda\cup\bigcup\limits_{j\neq j_0}S_j)$.

Claim 1. *The one-form ξ extends meromorphically to a neighborhood of q_{j_0} in M.*

We consider a local chart $(x,y)\in U$ for q_{j_0} as above. The restriction $\mathcal{F}|_U$ is transversely projective in $U\backslash\{y=0\}$. In particular, the holonomy map of the separatrix S_{j_0} for a simple loop $\gamma_{j_0}\subset S_{j_0}\backslash\{q_{j_0}\}$ around q_{j_0}, is of the form $h_{j_0}(u)=\frac{au}{1+bu}$ for some $a,b\in\mathbb{C}$, $a\neq0$ and some suitable holomorphic coordinate $u\in(\mathbb{C},0)$.

$1^{\mathbf{st}}$: $\mathcal{F}|_U$ is of type (Irred.i).

If $\lambda\notin\mathbb{Q}_-$ then $a=h'_{j_0}(0)=e^{2\pi i/\lambda}$ is not a root of the unity and therefore the homography $u\mapsto\frac{au}{1+bu}$ can be linearized by another homography. In particular h_{j_0} is analytically linearizable and therefore the singularity $q_{j_0}\in\operatorname{sing}(\mathcal{F})$ is analytically linearizable. Suppose now that $q_0\in\operatorname{sing}(\mathcal{F})$ is not analytically linearizable. Then we must have $\lambda=-\frac{k}{\ell}$ for some $\ell,k\in\mathbb{N}$, $\langle\ell,k\rangle=1$ and the holonomy h_{j_0} is analytically conjugate to the corresponding holonomy of the germ of singularity $\Omega_{k,\ell}=ky\,dx+\ell x(1+\frac{\sqrt{-1}}{2\pi}\,x^ky^\ell)dy$; such a singularity is called a *nonlinearizable resonant saddle.* Therefore, by [9] and [7] we may assume that $\mathcal{F}|_U$ is of the form $\Omega_{k,\ell}=0$ in the variables $(x,y)\in U$.

$2^{\mathbf{nd}}$: $\mathcal{F}|_U$ is of type (Irred.ii.a).

The singularity is a saddle-node with strong separatrix contained in $S_{j_0}:\{x=0\}$ and with holonomy of the form $h_{j_0}(u)=\frac{au}{1+bu}$ thus $a=1$, $k=1$ and h_{j_0} is conjugate to the holonomy of the strong separatrix of the germ of saddle-node singularity $\Omega_2=x^2\,dy-y\,dx$. This implies (cf. [8]) that $\mathcal{F}|_U$ can be given by $\Omega_2=0$ for suitable coordinates $(x,y)\in U$.

Summarizing we can assume that $\mathcal{F}|_U$ is of one of the following forms:

(a) $x\,dy-\lambda y\,dx=0$, $\lambda\neq0$

(b) $ky\,dx + \ell x(1 + \frac{\sqrt{-1}}{2\pi}\,x^k y^\ell)dy = 0, \quad k, \ell \in \mathbb{N}, \quad \langle k, \ell\rangle = 1.$
(c) $x^2 dy - y\,dx = 0$
with $\Lambda : \{y = 0\}$ and $S_{j_0} : \{x = 0\}$.

Case (a). In case (a) we write $\Omega(x, y) = g(x\,dy - \lambda y\,dx) = gx^{1+\lambda}\,d(yx^{-1})$ for some nonvanishing holomorphic function $g(x, y)$ and $\eta(x, y) = \frac{d(gx^{1+\lambda})}{gx^{1+\lambda}} + h\Omega = \frac{dg}{g} + (1+\lambda)\frac{dx}{x} + h\Omega$ for some holomorphic function $h(x, y)$. We define a local one-form ξ_0 in U by $\xi_0(x, y) := \frac{h^2\Omega}{2} - h\eta - dh$. Then ξ_0 is holomorphic and (Ω, η, ξ_0) is a projective triple in U. By Lemma 2 this implies that in $U^* = U\backslash\Lambda$ we have $\xi = \xi_0 + F\Omega$ where F is a holomorphic function satisfying (*) $d\Omega = -\frac{dF}{2F} \wedge \Omega$ in U^*. Now we show what this implies for F. We have from (*) that $d(g(xdy - \lambda ydx)) = -\frac{dF}{2F} \wedge (g(xdy - \lambda ydx))$ that is, $d(xdy - \lambda ydx) = -\frac{1}{2}\,\frac{d(Fg^2)}{Fg^2} \wedge (xdy - \lambda ydx)$. We set $G = Fg^2$ and write $G(x, y) = \sum\limits_{\substack{i\geq 0\\ j\in\mathbb{Z}}} G_{ij}\,x^i y^j$ in Laurent series. From the above we have $-2(1 + \lambda)G = x\,G_x + \lambda y G_y$ and therefore $-2(1 + \lambda) \cdot G_{ij} = (i + \lambda j)G_{ij}\,, \quad \forall\, i \geq 0, \forall\, j \in \mathbb{Z}$. If there is $G_{ij} \neq 0$ with $(i, j) \neq (0, 0)$ then $-2(1 + \lambda) = i + \lambda j$ and $-2 - i = \lambda(j + 2)$. Since $i \geq 0$ we have $2 + i \geq 2$ and therefore necessarily $\lambda = -\frac{(2+i)}{j+2} \in \mathbb{Q}$. But $\lambda \notin \mathbb{Q}_+$ thus $j + 2 \geq 1$ and therefore $j \geq -1$. We write therefore $\lambda = -\frac{k}{\ell}$ and obtain $\frac{k}{\ell} = \frac{2+i}{j+2} \quad \Rightarrow \quad k(j + 2) = \ell(2 + i) \Rightarrow \quad j = \frac{\ell(2+i)-2k}{k} = \frac{\ell(2+i)}{k} - 2$. Since $\langle \ell, k\rangle = 1$ we must have $2 + i = nk$ with $n \geq 1$ and therefore $j = n\ell - 2, \quad i = nk - 2$ with $i \geq 0, \quad j \geq -1$. This gives $G(x, y) = \sum\limits_{n=[\frac{2}{k}]}^{\infty} G_n \cdot x^{nk-2}\,y^{n\ell-2}$ for some coefficients G_n. We can rewrite $G(x, y) = \frac{1}{x^2y^2}(\sum_{n=[\frac{2}{k}]}^{+\infty} G_n \cdot (x^k y^\ell)^n) = \frac{\varphi(x^k y^\ell)}{x^2y^2}$ for a holomorphic function $\varphi(z) \in \mathbb{C}\{\{z\}\}$. If $\frac{2}{k} \leq 1$ then $\varphi(z) = \sum\limits_{n=1}^{\infty} G_n z^n$ and always G is holomorphic along $\{x = 0\}$. If $\frac{2}{k} > 1$ then $k = 1$ and $\frac{2}{k} = 2$ so that $\varphi(z) = \sum\limits_{n=2}^{+\infty} G_n z^n$. We conclude therefore that $G(x, y) = \frac{\varphi(x^k y^\ell)}{x^2y^2}$ where the holomorphic function $\varphi(z) \in \mathbb{C}\{\{z\}\}$ satisfies $\varphi(0) = 0$. Thus we have either $G(x, y)$ holomorphic in U or $G(x, y)$ has order one poles in $\{y = 0\}$ and is holomorphic in $U\backslash\{y = 0\}$. Since $G = Fg^2$ in $U^* = U\backslash\{y = 0\}$ this implies that $F = \frac{G}{g^2}$ extends meromorphically to U. This extension is either holomorphic or has poles of order one in $\{y = 0\}$ and is holomorphic in U^*. Assume now that $\lambda \notin \mathbb{Q}_-$ then $G_{ij} = 0, \ \forall (i, j) \neq (0, 0)$ and G is constant in U^*. This implies that F extends to U holomorphically as $F = \frac{G_{oo}}{g^2}\cdot$

Case (b). Now we consider case (b). Write $\Omega(x, y) = g\,\Omega_{k,\ell} = gx^{k+1}y^{\ell+1}\frac{1}{x^{k+1}y^{\ell+1}}[hydx + \ell x(1 + \frac{\sqrt{-1}}{2\pi}x^k y^\ell)dx] = gx^{k+1}y^{\ell+1}\,\omega_{k,\ell}$ where by definition $\omega_{k,\ell} = \frac{kdx}{y^\ell x^{k+1}} + \frac{\ell dy}{x^k y^{\ell+1}} + \frac{\sqrt{-1}}{2\pi}\,\ell\,\frac{dy}{y}$. We define $h(x, y)$ by $\eta(x, y) := \frac{d(gx^{k+1}y^{\ell+1})}{gx^{k+1}y^{\ell+1}} + h\Omega \Rightarrow \eta(x, y) = \frac{dg}{g} + (k + 1)\frac{cx}{x} + (\ell + 1)\frac{dy}{y} + h\Omega$ and ξ_0 by $\xi_0(x, y) := \frac{h^2\Omega}{2} - h\eta - dh$. Then $\xi = \xi_0 + F\Omega$ and as above $d\Omega = -\frac{dF}{2F} \wedge \Omega$ in U^*. Then from $\Omega = gx^{k+1}y^{\ell+1}\,\omega_{k,\ell}$ and from $d\omega_{k,\ell} = 0$ we obtain $d\omega_{k,\ell} = -\frac{dG}{2G} \wedge \omega_{k,\ell}$ in U^*. We have $\omega_{k,\ell} = df_{k,\ell}$ where $f_{k,\ell} = \frac{hx^k}{hx^k \log y+1}$ and $\omega_{k,\ell}$ (i.e., $\Omega_{k,\ell}$) admits no meromorphic first integral in U^*. Indeed, according to [7] $\Omega_{k,\ell}$ admits exactly two separatrices and is therefore nondicritical. If $\Omega_{k,\ell}$ admits a meromorphic first integral the, since it is nondicritical, it admits a holomorphic first integral. Since $\Omega_{k,\ell}$ is already irreducible it follows from [9] that $\Omega_{k,\ell} = 0$ is analytically linearizable, contradiction. Therefore, similarly to in the above argumentation we conclude that G is constant in U^* and therefore F extends holomorphically to U as $F = \frac{\text{cte.}}{g^2}\cdot$

Case (c). Finally, assume that we are in case (c). Write $\Omega(x, y) = g(x^2dy - ydx) = gx^2y(\frac{dy}{y} - \frac{dx}{x^2})$

then $\Omega(x,y) = gx^2y \cdot \omega(x,y)$ where $\omega(x,y) = \frac{dy}{y} - \frac{dx}{x^2} \cdot$ Again we introduce $\xi_0(x,y) = \frac{h^2\Omega}{2} - h\eta - dh$ and $G = Fg^2$ where $\xi = \xi_0 + F\Omega$ in U^*. We have $d\Omega = -\frac{dF}{2F} \wedge \Omega$ resulting in $d\omega = -\frac{dG}{2G} \wedge \omega$ in U^*. We have $\omega = d(\ell n\, y + \frac{1}{x}) = \frac{d\left(ye^{\frac{1}{x}}\right)}{ye^{\frac{1}{x}}}$ and it is well-known that a saddle-node singularity $\omega = 0$ (i.e., Ω) does not admit a nonconstant meromorphic first integral in U^* (see [8]). Again F extends holomorphically to U as $F = \frac{\text{cte.}}{g^2} \cdot$ This shows Claim 1 and indeed more. It is clear that (Ω, η, ξ) is a projective triple. We have proved that the projective triple (Ω, η, ξ) (as well as the projective transverse structure of $\mathcal{F}$) extends to a neighborhood of q_{j_0} in M. By Hartogs' Extension Theorem (see [6]) the one-form ξ extends to a neighborhood of $\Lambda \backslash \operatorname{sing}(\mathcal{F})$ in M and so does the projective transverse structure of $\mathcal{F}$. Therefore we have a projective triple (Ω, η, ξ) in $M \backslash \bigcup\limits_{j \neq j_0} S_j$. $\square$

4 Resolution of singularities and projective transverse structures

Now we consider a more general situation than the one in § 2. Let $\mathcal{F}$ be a holomorphic foliation with isolated singularities on a complex surface M^2, $\Lambda \subset M$ an invariant compact analytic subset of dimension one and such that $\operatorname{sing}(\mathcal{F}) \subset \Lambda$. We perform the Resolution of Singularities for $\mathcal{F}$ in M which gives (cf. [12], [2], [9]): a complex surface $\widetilde{M}^2$ and a proper holomorphic map $\pi \colon \widetilde{M} \to M$ which is a finite composition of quadratic blow-ups such that $P = \pi^{-1}(\Lambda)$ is a normal crossing divisor without triple points $P = \widetilde{\Lambda} \cup E$ where $\widetilde{\Lambda} = \overline{\pi^{-1}(\Lambda \backslash \operatorname{sing}(\mathcal{F}))}$, $E = \pi^{-1}(\operatorname{sing}(\mathcal{F}))$ is a union of projective lines $E = \bigcup\limits_{j=1}^{m} E_j$, $E_j \simeq \mathbb{C}P(1)$ with negative self-intersection in $\widetilde{M}$, the pull-back $\widetilde{\mathcal{F}} = \pi^*(\mathcal{F})$ is a foliation with isolated singularities, $\operatorname{sing}(\widetilde{\mathcal{F}}) \subset \pi^{-1}(\operatorname{sing}(\mathcal{F}))$ consists of irreducible singularities of types (Irred.i), (Irred.ii.a) and (Irred.ii.b) as in § 3 above. Any component $E_j \subset E$ is either invariant or everywhere transverse to $\widetilde{\mathcal{F}}$ (in this last case we shall say that E_j is a *dicritical component*). Write $\Lambda = \bigcup\limits_{j=1}^{r} \Lambda_j$ in (compact) irreducible components in M. Then we can write $\widetilde{\Lambda} = \bigcup\limits_{j=1}^{r} \widetilde{\Lambda}_j$ in compact (invariant by $\widetilde{\mathcal{F}}$) irreducible components in $\widetilde{M}$, with $\widetilde{\Lambda}_j = \overline{\pi^{-1}(\Lambda_j \backslash \operatorname{sing}(\mathcal{F}))}$. We can also assume that if $\tilde{q} \in \operatorname{sing}(\widetilde{\mathcal{F}}) \subset P$ is a singularity of $\widetilde{\mathcal{F}}$ then the local separatrices of $\widetilde{\mathcal{F}}$ through $\tilde{q}$ are contained in P. This is not necessarily true for the inverse image of the separatrices of points in $\operatorname{sing}(\mathcal{F})$. Now we suppose that Ω is a meromorphic one-form defining $\mathcal{F}$ in M having polar set in general position with Λ. Also let η be a logarithmic derivative of Ω adapted to Λ. Then the forms $\widetilde{\Omega} = \pi^*(\Omega)$ and $\tilde{\eta} = \pi^*(\eta)$ have similar properties with respect to $\widetilde{\Lambda}$ and P (see [3]). Write $\operatorname{sep}(\mathcal{F}, \Lambda)$ for the set of separatrices of $\mathcal{F}$ not contained in Λ, this set may have infinitely many components. We have $\operatorname{sep}(\mathcal{F}, \Lambda) \cap \Lambda \subset \operatorname{sing}(\mathcal{F})$. Assume that $\mathcal{F}$ is transversely projective in $M \backslash [\Lambda \cup (\operatorname{sep}(\mathcal{F}, \Lambda) \backslash \bigcup\limits_{j \in A} S_j)]$ where $A = \{1, \ldots, t\}$ is a finite set of indices and each $S_j \subset \operatorname{sep}(\mathcal{F}, \Lambda)$ is a local separatrix. Then $\widetilde{\mathcal{F}}$ is transversely projective in $\widetilde{M} \backslash \pi^{-1}(\Lambda \cup (\operatorname{sep}(\mathcal{F}, \Lambda) \backslash \bigcup\limits_{j \in A} S_j)))$. However, we can say more. Given an invariant subset $\Gamma \subset P$ we denote by $\operatorname{sep}(\widetilde{\mathcal{F}}, \Gamma)$ the set of separatrices of $\widetilde{\mathcal{F}}$ (through singular points in Γ and) not contained in Γ. Let $j \in A$ and $\{q_j\} = S_j \cap \Lambda \subset \operatorname{sing}(\mathcal{F})$. We shall say that q_j is a *nondicritical* singularity to indicate that $\pi^{-1}(q_j)$ is invariant by $\widetilde{\mathcal{F}}$ and it is a union of projective lines. Put $E(q_j) = \pi^{-1}(q_j)$ and $\widetilde{S}_j := \overline{\pi^{-1}(S_j \backslash \{q_j\})} \subset \widetilde{M}$. We have $\widetilde{S}_j \subset P$ because q_j is nondicritical. By the choice of A and $\widetilde{S}_j$, $\widetilde{\mathcal{F}}$ is transversely projective in $\widetilde{M} \backslash (P \backslash \bigcup\limits_{j \in A} S_j)$. Let $\Gamma_j \subset E(q_j)$ be an irreducible component of $E(q_j)$ with $E(q_j) \cap \widetilde{S}_j = \Gamma_j \cap \widetilde{S}_j = \{\tilde{q}_j\} \subset \operatorname{sing}(\widetilde{\mathcal{F}})$ and assume that $\widetilde{S}_j$ *is not a central manifold*: this means that either $\tilde{q}_j$ is of type (Irred.i) or it is of type (II.i.a), i.e. a saddle-node singularity whose strong manifold is contained in $\widetilde{S}_j$. We denote by $\operatorname{Inv}(\widetilde{\mathcal{F}}) \subset P$ the invariant part of P and given any point $\tilde{p} \in \operatorname{Inv}(\widetilde{\mathcal{F}})$ by $\operatorname{Inv}(\tilde{p})$ the connected component of $\operatorname{Inv}(\widetilde{\mathcal{F}})$ that contains the point $\tilde{p}$. Thus, the singularity q_j is nondicritical if and only if $\operatorname{Inv}(\tilde{q}_j) = E(q_j)$. Notice that $\widetilde{S}_j \subset \operatorname{sep}(\widetilde{\mathcal{F}}, \Gamma_j)$ and that there is an open neighborhood W_j of Γ_j in $\widetilde{M}$ such that $\widetilde{\mathcal{F}}$ is transversely projective in $W_j \backslash [\Gamma_j \cup (\operatorname{sep}(\widetilde{\mathcal{F}}, \Gamma_j) \backslash \widetilde{S}_j)]$. Therefore, applying Lemma 3 above we conclude that $\widetilde{\mathcal{F}}$ admits a projective triple $(\widetilde{\Omega}, \widetilde{\eta}, \widetilde{\xi})$ in $\widetilde{M} \backslash P$ and $\widetilde{\xi}$ extends giving a projective triple in the neighborhood W_j of Γ_j in $\widetilde{M}$. Let us take any other component Γ_j' of $E(q_j)$ *adjacent* to Γ_j, i.e., $\Gamma_j' \cap \Gamma_j \neq \emptyset$. If Γ_j' is not $\widetilde{\mathcal{F}}$-invariant then clearly the projective transverse structure as well as the projective triple $(\widetilde{\Omega}, \widetilde{\eta}, \widetilde{\xi})$ extend to a neighborhood of $\Gamma_j' \backslash (\operatorname{Inv}(\mathcal{F}) \cap \Gamma_j')$. Hence we suppose that Γ_j' is also $\widetilde{\mathcal{F}}$-invariant. In particular we have a corner singularity $\{\tilde{q}_j\} = \Gamma_j \cap \Gamma_j'$, $\tilde{q}_j \in \operatorname{sing}(\widetilde{\mathcal{F}})$. If $\tilde{q}_j$ is such that Γ_j' is not a central manifold for $\tilde{q}_j$ then Lemma 3 applies to show that the projective triple $(\widetilde{\Omega}, \widetilde{\eta}, \widetilde{\xi})$ extends with a projective transverse structure for $\widetilde{\mathcal{F}}$ in a neighborhood of $\tilde{q}_j$. Put $\mathcal{S} = \bigcup\limits_{j \in A} S_j \subset \operatorname{sep}(\mathcal{F}, \Lambda)$.



Figure 3: Arrows diagram for irreducible singularities (Irred.i), (Irred.ii.a) and $\Gamma_{\mathcal{S}}(\tilde{q})$.

Summarizing the above discussion we have (see Figure 3):

Definition 2. Given a singularity $q \in \operatorname{sing}(\mathcal{F})$ with $q \in \mathcal{S}$ and any singularity $\tilde{q} \in E(q)$ denote by $\Gamma_{\mathcal{S}}(\tilde{q})$ the maximal subset of $\operatorname{Inv}(\tilde{q}) \subset \operatorname{Inv}(\widetilde{\mathcal{F}})$ such that:

(i) There exists some separatrix $\widetilde{S}_j$, $j \in A$ with $\widetilde{S}_j \cap \Gamma_{\mathcal{S}}(\tilde{q}) \neq \emptyset$ and which is not a central manifold.

(ii) If $\tilde{q}_0 \in \operatorname{Inv}(\tilde{q}) \cap \operatorname{sing}(\widetilde{\mathcal{F}})$ is a saddle-node singularity then $\Gamma_{\mathcal{S}}(\tilde{q})$ is not a central manifold for $\tilde{q}_0$.

Lemma 4. *The one-form $\widetilde{\xi}$ extends to a neighborhood of each $\Gamma_{\mathcal{S}}(\tilde{q})$ with a projective transverse structure in this neighborhood.*

Proposition 2. *Let $\mathcal{F}$, M, Λ, Ω, η, $\operatorname{sep}(\mathcal{F},\Lambda), S_j \subset \operatorname{sep}(\mathcal{F})$, $j \in A$ and $\mathcal{S} = \bigcup\limits_{j\in A} S_j$ as above. Suppose that:*
(i) *$\mathcal{F}$ is transversely projective in* $M\backslash[\Lambda \cup (\operatorname{sep}(\mathcal{F},\Lambda)\backslash\mathcal{S})] = M\backslash(\Lambda \cup [\operatorname{sep}(\mathcal{F},\Lambda)\backslash(\bigcup\limits_{j\in A} S_j)])$;
(ii) $\operatorname{Inv}(\widetilde{\mathcal{F}}) = \bigcup\limits_{\substack{\tilde{q}\in\operatorname{sing}(\widetilde{\mathcal{F}})\\ \pi(\tilde{q})\in\mathcal{S}}} \Gamma_{\mathcal{S}}(\tilde{q})$.
Then $\mathcal{F}$ admits a projective triple (Ω,η,ξ) with ξ transversely projective in a neighborhood of Λ in M.

Proof: By the above Lemma 4 there is a projective triple $(\widetilde{\Omega},\widetilde{\eta},\widetilde{\xi})$ in $\widetilde{M}$ where $\widetilde{\xi}$ is transversely projective in $\widetilde{M}$. Using the map $\pi\colon \widetilde{M} \to M$ to project $\widetilde{\xi}$ we obtain a one-form ξ in M with the desired properties. □

Definition 3. Let $\mathcal{F}$ be a holomorphic foliation with isolated singularities or a complex surface M^2 and $\Lambda \subset M^2$ a compact analytic invariant subset of dimension one. We shall say that $\mathcal{F}$ *has noncentral separatrices transverse to* Λ if there is a subset $\mathcal{S} = \bigcup\limits_{j\in A} S_j \subset \operatorname{sep}(\mathcal{F},\Lambda)$ as above such that $\operatorname{Inv}(\widetilde{\mathcal{F}}) = \bigcup\limits_{\substack{\tilde{q}\in\operatorname{sing}(\widetilde{\mathcal{F}})\\ \pi(\tilde{q})\in\mathcal{S}}} \Gamma_{\mathcal{S}}(\tilde{q})$.

5 Transversely projective foliations on the complex projective plane

In this section we study (transversely projective) foliations on the complex projective plane $\mathbb{C}P(2)$ and prove Theorems 1 and 3. Let $\mathcal{F}$ be a holomorphic foliation of codimension one on $\mathbb{C}P(2)$ having singular set $\operatorname{sing}(\mathcal{F}) \subsetneqq \mathbb{C}P(2)$. As it is well-known we can assume that $\operatorname{sing}(\mathcal{F})$ is of codimension ≥ 2 and $\mathcal{F}$ is given in any affine space $\mathbb{C}^2 \subset \mathbb{C}P(2)$ with coordinates (x,y), by a polynomial one-form $\Omega(x,y) = A(x,y)dx + B(x,y)dy$ with $\operatorname{sing}(\mathcal{F}) \cap \mathbb{C}^2 = \operatorname{sing}(\Omega)$. In particular $\operatorname{sing}(\mathcal{F}) \subset \mathbb{C}P(2)$ is a nonempty finite set of points. Given any algebraic subset $\Lambda \subset \mathbb{C}P(2)$ of dimension one we can therefore always obtain a meromorphic (rational) one-form Ω on $\mathbb{C}P(2)$ such that Ω defines $\mathcal{F}$, $(\Omega)_\infty$ is noninvariant and in general position in the sense of § 2 (indeed, we can assume that $(\Omega)_\infty$ is any projective line in $\mathbb{C}P(2)$). Also if we take $\eta_0 = \frac{B_x}{B}\,dx + \frac{A_y}{A}\,dy$ then we obtain a rational one-form such that $d\Omega = \eta_0 \wedge \Omega$ and with polar set given by $(\eta_0)_\infty = \{(x,y) \in \mathbb{C}^2 : A(x,y) = 0\} \cup \{(x,y) \in \mathbb{C}^2 : B(x,y) = 0\} \cup (\Omega)_\infty$. In particular, $(\eta_0)_\infty \cap \mathbb{C}^2$ has order one and the "residue" of η along any component C of $(\Omega)_\infty$ equals $-k$ where k is the order of C as a pole of Ω. Any rational one-form η such that $d\Omega = \eta \wedge \Omega$ writes $\eta = \eta_0 + h\Omega$ for some rational function h. We obtain in this way rational logarithmic derivatives η of Ω adapted to (Ω,Λ). Applying the results of § 2 we promptly obtain.

Proposition 3. *Let $\mathcal{F}$ be a holomorphic foliation on $\mathbb{C}P(2)$. Assume that $\mathcal{F}$ is transversely projective in $\mathbb{C}P(2)\backslash\Lambda$ for some algebraic subset Λ of dimension one. Then $\mathcal{F}$ has a projective triple (Ω,η,ξ) where Ω and η are rational one-forms and ξ is meromorphic on $\mathbb{C}P(2)\backslash\Lambda$. In particular ξ defines a transverse foliation $\mathcal{F}^\perp$ to $\mathcal{F}$ on $\mathbb{C}P(2)\backslash\Lambda$ having a projective transverse structure.*

Proof of Theorem 3. According to Proposition 3 we have a projective triple (Ω,η,ξ) where Ω, η are rational one-forms and ξ is meromorphic in $\mathbb{C}P(2)\backslash\Lambda$. By hypothesis $\mathcal{F}$ has noncentral separatrices transverse to Λ therefore according to Proposition 2 the one-form ξ extends to a meromorphic one-form on $\mathbb{C}P(2)$ and the corresponding transverse foliation $\mathcal{F}^\perp$ defined by ξ is also transversely projective in a neighborhood of Λ on $\mathbb{C}P(2)$. Thus $\mathcal{F}^\perp$ is transversely projective on $\mathbb{C}P(2)$. This implies that $\mathcal{F}^\perp$ admits a rational first integral $R\colon \mathbb{C}P(2) \dashrightarrow \overline{\mathbb{C}}$ ([10]). If we write $\xi = g\,dR$ for some rational function g then we may replace the rational triple (Ω, η, ξ) by (Ω', η', ξ') where $\Omega' = g\Omega$, $\eta' = \eta + \frac{dg}{g}$ and $\xi' = \frac{1}{g}\,\xi = dR$. The relations $d\Omega' = \eta' \wedge \xi'$, $d\eta' = \Omega' \wedge \xi'$, $d\xi' = \xi \wedge \eta'$ imply that $\eta' = HdR$ for some rational function H. Now we define $\omega := \frac{H^2}{2}\,\xi' - H\eta' + dH = \frac{1}{2}\,H^2 dR + dH$ one-form such that $d\omega = -HdH \wedge dR$. On the other hand $\eta' \wedge \omega = HdR \wedge dH = -HdH \wedge dR$. Thus $d\omega = \eta' \wedge \omega$. We also have $d\eta' = dH \wedge dR = (-\frac{1}{2}\,H^2 dR + dH) \wedge dR = \omega \wedge \xi'$. The rational triple (ω, η', ξ') satisfies the projective relations $d\omega = \eta' \wedge \omega$, $d\eta' = \omega \wedge \xi'$, $d\xi' = \xi' \wedge \eta'$ and therefore by a natural adaptation of Lemma 2 we conclude that $\Omega' = \omega + F.\xi'$ for some rational function ℓ such that $d\xi' = \xi' \wedge \frac{dF}{2F}\cdot$ This implies $dF \wedge dR \equiv 0$. By Stein Fatorization Theorem we may assume from the beginning that R has connected fibers and therefore $dF \wedge dR \equiv 0$ implies $\ell = \varphi(R)$ for some one-variable rational function $\varphi(z) \in \mathbb{C}(z)$. We obtain therefore $\Omega' = -\frac{1}{2}\,H^2 dR + dH + \varphi(R)dR == dH - (\frac{1}{2}\,H^2 - \varphi(R))dR$. If we define a rational map $\sigma\colon \mathbb{C}P(2) \dashrightarrow \overline{\mathbb{C}} \times \overline{\mathbb{C}}$ by $\sigma(x,y) = \big(R(x,y), H(x,y)\big)$ on $\mathbb{C}^2$ then clearly $\Omega' = \sigma^*(dy - (\frac{1}{2}\,y^2 - \varphi(x))dx)$ and therefore $\mathcal{F}$ is the pull-back $\mathcal{F} = \sigma^*(\mathcal{R})$ of the Riccati foliation $\mathcal{R}$ given on $\overline{\mathbb{C}} \times \overline{\mathbb{C}}$ by the rational one-form $dy - (\frac{1}{2}\,y^2 - \varphi(x))dx$. □

Proof of Theorem 1. First of all we make a remark. We can assume that Λ is minimal with respect to the projective structure of the transverse foliation $\mathcal{F}^\perp$ given in Proposition 3, in the sense that there is no proper algebraic invariant curve $\Lambda^* \subsetneqq \Lambda$ such that $\mathcal{F}^\perp$ is defined and transversely projective in the complement of Λ^*. If $\Lambda = \emptyset$ then $\mathcal{F}^\perp$ is transversely projective in $\overline{\mathbb{C}} \times \overline{\mathbb{C}}$ and therefore it admits a rational first integral. This implies, as in the proof of Theorem 3, that $\mathcal{F}$ is a rational pull-back of a Riccati foliation. Thus we assume that Λ is minimal and nonempty. Given a singularity $q \in \operatorname{sing}(\mathcal{F})$ such that $q \notin \Lambda$ then $\mathcal{F}$ is transversely projective in a punctured neighborhood of q and therefore $\mathcal{F}$ admits a meromorphic first integral in a neighborhood of q ([10]). Hence such a singularity cannot be hyperbolic. Since by hypothesis all singularities of $\mathcal{F}$ are hyperbolic we conclude that $\operatorname{sing}(\mathcal{F}) \subset \Lambda$. Given a hyperbolic singularity $q \in \operatorname{sing}(\mathcal{F}) \cap \Lambda$ there are local coordinates $(x,y) \in U$ centered at q such that $\mathcal{F}\big|_U$ is given by $xdy - \lambda ydx = 0$, $\lambda \in \mathbb{C} \setminus \mathbb{R}$. In particular $\mathcal{F}$ has exactly two separatrices $\{x = 0\}$ and $\{y = 0\}$ through q. We may assume that $\{y = 0\} \subset \Lambda$. Suppose that $\{x = 0\} \not\subset \Lambda$ then $\{x = 0\}$ is a noncentral separatrix transverse to Λ as in Theorem 3. According to Lemma 3 and also to § 4 the transverse foliation $\mathcal{F}^\perp$ and its projective transverse structure extend to a neighborhood of $(\Lambda_1 \setminus (\Lambda_1 \cap \operatorname{sing}(\mathcal{F}))) \cup \{q\}$ where Λ_1 is the irreducible component of Λ that contains the local separatrix $\{y = 0\}$. Thus we conclude that $\mathcal{F}^\perp$ is defined and transversely projective in the complement of the algebraic curve Λ^* obtained by deleting Λ_1 in Λ. Since we are assuming that Λ is minimal we obtain a contradiction. Therefore, we conclude that $\{x.y = 0\} \subset \Lambda$, i.e., Λ contains all the separatrices through the singularities $q \in \operatorname{sing}(\mathcal{F})$. Thus, Λ is an invariant algebraic curve which has normal crossings and passes through all the singularities of $\mathcal{F}$ and contains all the separatrices of the singularities of $\mathcal{F}$. According to [1] this implies that $\mathcal{F}$ is a logarithmic foliation. □

6 The case of dimension $m \geq 3$

In this section $\mathcal{F}$ denotes a holomorphic foliation of codimension one on $\mathbb{C}P(m)$, $m \geq 3$, having singular set $\operatorname{sing}(\mathcal{F})$ of codimension ≥ 2. We assume that $\mathcal{F}$ is transversely projective in $\mathbb{C}P(m) \setminus \Lambda$ for some codimension one invariant algebraic subset $\Lambda \subset \mathbb{C}P(m)$. A very basic remark is (cf. [10]): let $\sigma\colon \mathbb{C}P(2) \to \mathbb{C}P(m)$ be a linear embedding and assume that σ is in *general position* with respect to $\mathcal{F}$ and Λ, i.e., σ is transverse to $\Lambda \cup \operatorname{sing}(\mathcal{F})$ and the foliation $\mathcal{F}^* = \sigma^*(\mathcal{F})$ has isolated singularities on $\mathbb{C}P(2)$. Any linear embedding $\mathbb{C}P(2) \to \mathbb{C}P(m)$ can be arbitrarily approximated by one in general position as above.

Lemma 5. *If σ is general position then $\mathcal{F}^*$ is transversely projective outside the invariant algebraic curve $\Lambda^* = \sigma^{-1}(\Lambda) = \Lambda \cap \sigma(\mathbb{C}P(2))$.*

We shall refer to $\mathcal{F}^*$ as a *generic* (*linear*) *section of* $\mathcal{F}$ and we have

Proposition 4. *Let $\mathcal{F}^*$ be a generic linear section of $\mathcal{F}$. Suppose that $\mathcal{F}^*$ is a rational pull-back of a Riccati foliation and that $\mathcal{F}$ is transversely projective in $\mathbb{C}P(m)\backslash\Lambda$. Then $\mathcal{F}$ is a rational pull-back of a Riccati foliation.*

We shall need the following:

Lemma 6. *Let $\mathcal{F}$ be a codimension one holomorphic foliation on $\mathbb{C}P(m)$, $m \geq 3$. Then there is a pair Ω, η of rational one-forms such that $\mathcal{F}$ is defined by Ω and η is a logarithmic derivative of Ω.*

Proof: It is well-known that given any affine space $\mathbb{C}^m \subset \mathbb{C}P(m)$ with affine coordinates $(x_1, \ldots, x_m)$ there is a polynomial one-form $\Omega = \sum\limits_{j=1}^{m} P_j\, dx_j$ which defines $\mathcal{F}|_{\mathbb{C}^m}$ and with $g.c.d.(P_1, \ldots, P_m) = 1$; also Ω must satisfy the integrability condition $\Omega \wedge d\Omega = 0$. Hence $0 = \Omega \wedge d\Omega = (\sum\limits_{j=1}^{m} P_j dx_j) \wedge (\sum\limits_{i=1}^{m} \sum\limits_{k=1}^{m} \frac{\partial P_i}{\partial x_k}\, dx_k \wedge dx_i)$. This gives $\sum\limits_{i,j,k=1}^{m} \frac{\partial P_i}{\partial x_k} \cdot P_j\, dx_j \wedge dx_k \wedge dx_i = 0$ and therefore

$$0 = \sum_{i<j<k} [(\frac{\partial P_i}{\partial x_k} \cdot P_j - \frac{\partial P_i}{\partial x_j} \cdot P_k) + (\frac{\partial P_j}{\partial x_k} \cdot P_i - \frac{\partial P_j}{\partial x_i} \cdot P_k)] dx_j dx_k dx_i \qquad (*)$$

Now define in a natural way $\eta := \sum_{j=1}^{m} (\sum_{k \neq j} \frac{1}{P_k} \cdot \frac{\partial P_k}{\partial x_j}) dx_j$. From (*) we obtain $d\Omega = \eta \wedge \Omega$. $\square$

Proof of Proposition 4: We identify $\mathbb{C}P(2)$ with $\sigma(\mathbb{C}P(2))$. Assume first that $m = 3$. Put $\Omega^* = \sigma^*(\Omega)$, $\eta^* = \sigma^*(\eta)$ and let ξ^* be a rational one-form on $\mathbb{C}P(2)$ such that $(\Omega^*, \eta^*, \xi^*)$ is projective and ξ^* has a rational first integral on $\mathbb{C}P(2)$: the existence of ξ^* as above is a consequence of the fact that $\mathcal{F}^*$ is a rational pull-back of a Riccati foliation. Locally, outside the singular set of $\mathcal{F}^*$, we can write $\xi^* = g_\alpha df_\alpha$, $\eta^* = \frac{dg_\alpha}{g_\alpha} + h_\alpha\, g_\alpha\, df_\alpha$ and $\Omega^* = \frac{h_\alpha^2}{2}\, \xi^* - h_\alpha \eta^* + dh_\alpha = \frac{h_\alpha^2}{2}\, g_\alpha\, df_\alpha - h_\alpha(\frac{dg_\alpha}{g_\alpha} + h_\alpha\, g_\alpha\, df_\alpha) + dh_\alpha = -\frac{h_\alpha^2}{2}\, g_\alpha\, df_\alpha - h_\alpha\, \frac{dg_\alpha}{g_\alpha} + dh_\alpha$, for some holomorphic functions f_α , g_α , h_α in open subsets U_α which cover $\mathbb{C}P(2) \backslash \operatorname{sing}(\mathcal{F})$. By transversality we can extend locally f_α , g_α and h_α as constant along the leaves of $\mathcal{F}$ by using local flow boxes for $\mathcal{F}$. This process gives local extensions ξ_α of ξ^* as well as local extensions η_α and Ω_α of η^* and Ω^* to open subsets V_α on $\mathbb{C}P(m)$ such that $V_\alpha \cap \mathbb{C}P(2) = U_\alpha$. Nevertheless we have global one-forms Ω and η with $\Omega|_{V_\alpha} = \Omega_\alpha$ and $\eta|_{V_\alpha} = \eta_\alpha$. Therefore, associated to an intersection $U_\alpha \cap U_\beta \neq \emptyset$ we must have $V_\alpha \cap V_\beta \neq \emptyset$ and in $V_\alpha \cap V_\beta$: $\xi_\alpha = \xi_\beta + F_{\alpha\beta}\Omega$ for some meromorphic function $F_{\alpha\beta}$ such that $d\Omega = -\frac{dF_{\alpha\beta}}{2F_{\alpha\beta}} \wedge \Omega$ and $F_{\alpha\beta}|_{U_\alpha \cap U_\beta} \equiv 0$. Since $F_{\alpha\beta}$ is obtained from $F^*_{\alpha\beta} = F_{\alpha\beta}|_{U_\alpha \cap U_\beta}$ by trivial extension via flow box for $\mathcal{F}$ we conclude that $F_{\alpha\beta} \equiv 0$ and therefore $\xi_\alpha = \xi_\beta$ in $V_\alpha \cap V_\beta$. This shows that we have a coherent extension of ξ^* to a neighborhood of $\mathbb{C}P(2) \backslash \operatorname{sing}(\mathcal{F})$ and by Hartogs' Extension Theorem we conclude that ξ^* admits an extension to a neighborhood of $\mathbb{C}P(2)$ on $\mathbb{C}P(3)$ and there since $\mathbb{C}P(3) \backslash \mathbb{C}P(2)$ is a Stein space we extend ξ^* to $\mathbb{C}P(3)$, denote by ξ this rational one-form on $\mathbb{C}P(3)$. By the Identity Principle ξ also defines a projective triple (Ω, η, ξ). Since the restriction $\xi^* = \xi|_{\mathbb{C}P(2)}$ admits a rational first integral the one-form ξ admits a rational first integral on $\mathbb{C}P(3)$. Using [10] (cf. Theorem 4.1 § 4) or the same arguments in the end of the proof of Theorem 1 we conclude that $\mathcal{F}$ is a rational pull-back of a Riccati foliation. This proves the case $m = 3$. Proceeding in this way by induction we prove the general case $m \geq 3$. $\square$

Now it is easy to deduce the following codimension one version of Theorem 3:

Theorem 4. *Let $\mathcal{F}$ be a codimension one holomorphic foliation on $\mathbb{C}P(m)$, $m \geq 2$ such that $\mathcal{F}$ is transversely projective outside an algebraic invariant subset $\Lambda \subset \mathbb{C}P(m)$ of codimension one. Assume that a generic linear section $\mathcal{F}^*$ of $\mathcal{F}$ has noncentral separatrices transverse to Λ^*. Then $\mathcal{F}$ is a rational pull-back of a Riccati foliation.*

7 Foliations on compact manifolds

In this section we apply some results of [4] together with our study in order to give a description of the "generic case" for a foliation on a compact manifold of dimension ≥ 3. Let $\mathcal{F}$ be a codimension one holomorphic foliation with singular set $\operatorname{sing}(\mathcal{F})$ of codimension ≥ 2 on a compact complex manifold M^m of dimension $m \geq 3$. We have the following Algebraic Reduction Theorem:

Theorem 5 ([4]). *Let $\sigma\colon M \to \mathcal{A}(M)$ be an algebraic reduction of M and assume that $\mathcal{F}$ is pseudo-parallelizable. Then either $\mathcal{F}$ is the pull-back by a meromorphic map of an algebraic foliation $\mathcal{F}_1$ on the manifold $\mathcal{A}(M)$ or $\mathcal{F}$ admits a meromorphic triple (Ω, η, ξ) on M where $\mathcal{F}$ is given by Ω and we have the projective relations $d\Omega = \eta \wedge \Omega$, $d\eta = \Omega \wedge \xi$ and $d\xi = \xi \wedge \eta$.*

The notion of *pseudo-parallelizable* is verified if there is a meromorphic vector field X on M which is generically transverse to $\mathcal{F}$. The foliation $\mathcal{F}$ is a meromorphic pull-back of a foliation on a lower dimensional manifold in case the *algebraic dimension* of M is not equal to $\dim(M)$. In this case the pull-back is the composition of the algebraic reduction of M with a sequence of blow-ups at smooth centers. Motivated by this theorem we shall say that a codimension one holomorphic foliation $\mathcal{F}$ on a compact manifold M^m having singular set $\operatorname{sing}(\mathcal{F})$ of codimension ≥ 2, is *pseudo-algebraic* if there is a meromorphic map $\tau\colon M \to N$ from M into an algebraic projective manifold N and an algebraic foliation $\mathcal{F}_1$ of codimension one on N such that $\mathcal{F} = \tau^*(\mathcal{F}_1)$. Notice that we do not require the dimension of N to be smaller than the dimension of M. Nevertheless, if $\dim N \geq \dim M$ then necessarily M is algebraic so the notion is not interesting in this case.

Now we can prove Theorem 2.

Proof of Theorem 2: Assume that $\mathcal{F}$ is not pseudo-algebraic. Then by Theorem 5 there is a meromorphic triple of one-forms Ω, η, ξ on M such that Ω defines $\mathcal{F}$ and we have the projective relations $d\Omega = \eta \wedge \Omega$, $d\eta = \Omega \wedge \xi$, $d\xi = \xi \wedge \eta$ on M. It follows that $\mathcal{F}$ is transversely projective in $M \backslash \Lambda$ for some compact analytic subset $\Lambda \subset M$ of codimension one (see § 2). We have $\Lambda \subset (\Omega)_\infty \cup (\eta)_\infty \cup (\xi)_\infty$. If η can be chosen to be closed then we are in case (ii). We may assume therefore that η is not closed, ξ not identically zero and Λ is minimal with the property that $\mathcal{F}$ is transversely projective in $M \backslash \Lambda$. If $\Lambda = \emptyset$ then $\mathcal{F}$ is transversely projective in M and since M is simply-connected $\mathcal{F}$ admits a meromorphic first integral on M. We can suppose $\Lambda \neq \emptyset$. Suppose now that for some irreducible component Λ_1 of Λ we have $\operatorname{sep}(\mathcal{F}, \Lambda_1) \neq \emptyset$; then there is some separatrix $S_1 \in \operatorname{sep}(\mathcal{F}, \Lambda_1)$ transverse to Λ_1 with $S_1 \cap \Lambda_1 \subset \operatorname{sing}_2(\mathcal{F})$ the codimension two component of $\operatorname{sing}(\mathcal{F})$. The same argumentation in the Extension Lemma 3 in § 2 shows that the projective transverse structure of ξ extends to a neighborhood of Λ_1 (notice that there are no singularities like saddle-nodes or resonant saddles in Λ_1). Thus ξ is transversely projective in M and since M is simply-connected we obtain a meromorphic first integral for ξ on M. Using [10] or the proof of Theorem 1 (adapted to this case) we conclude that $\mathcal{F}$ is a meromorphic pull-back of a Riccati foliation. The only remaining case is as follows. There is a codimension one compact analytic subset $\Lambda \subset M$ invariant by $\mathcal{F}$ such that $\mathcal{F}$ is transversely projective on $M \backslash \Lambda$ and for any irreducible component Λ_1 of Λ we have $\operatorname{sep}(\mathcal{F}, \Lambda_1) = \emptyset$, i.e., Λ_1 contains all separatrices through the points in $\operatorname{sing}(\mathcal{F}) \cap \Lambda_1$. Notice that a singularity in $\operatorname{sing}_2(\mathcal{F})$ is by hypothesis hyperbolic.

Therefore $\mathcal{F}$ cannot be transversely projective in any neighborhood of this singularity because this would imply the existence of a meromorphic first integral in a neighborhood of this point, contradiction. Therefore we must have $\operatorname{sing}_2(\mathcal{F}) \subset \Lambda$. Now, applying arguments similar to the ones in the proof of Theorem 1 we conclude that $\mathcal{F}$ is a logarithmic foliation. $\square$

References

[1] M. Brunella; *Birrational geometry of foliations.* Lecture Notes of the First Latin American Congress of Mathematicians, IMPA - Rio de Janeiro (2000).

[2] C. Camacho, P. Sad; *Invariant varieties through singularities of holomorphic vector fields.* Ann. of Math. 115 (1982), 579–595.

[3] C. Camacho, B. Scárdua; *Holomorphic foliations with Liouvillian first integrals.* Ergodic Theory and Dynamical Systems (2001), 21, pp.717-756.

[4] D. Cerveau. A. Lins Neto, F. Loray, J. Pereira, F. Touzet; *Algebraic reduction for holomorphic foliations.* Pre-publ. IRMAR - Rennes, 2004.

[5] C. Godbillon; Feuilletages: Études geométriques. Birkhäuser.

[6] R. C. Gunning; Introduction to holomorphic functions of several variables. Vol. II. Local theory. The Wadsworth & Brooks/Cole Mathematics Series. Wadsworth & Brooks/Cole Advanced Books & Software, Monterey, CA, 1990.

[7] J. Martinet et J-P. Ramis; *Classification analytique des équations différentielles nonlineaires resonnants du premier ordre.* Ann. Sc. Ec. Norm. Sup., 16,1983, 571-621.

[8] J. Martinet et J-P. Ramis; *Problème de modules pour des équations différentielles non lineaires du premier ordre.* Publ. Math. Inst. Hautes Études Scientifiques, 55 (1982), 63-124.

[9] J.F. Mattei and R. Moussu; *Holonomie et intégrales premières.* Ann. Ec. Norm. Sup. 13 (1980), 469–523.

[10] B. Azevedo Scárdua; *Transversely affine and transversely projective foliations.* Ann. Sc. École Norm. Sup., 4^e série, t.**30**, 1997, p.169-204

[11] B. Azevedo Scárdua; *Integration of complex differential equations.* Journal of Dynamical and Control Systems, issue 1, vol. **5**, pp.1-50, 1999.

[12] A. Seidenberg; *Reduction of singularities of the differential equation $Ady = Bdx$.* Amer. J. of Math. 90 (1968), 248–269.

Bruno Scardua
Instituto de Matemática
Universidade Federal do Rio de Janeiro
Caixa Postal 68530
CEP. 21945-970 Rio de Janeiro - RJ
BRAZIL

scardua@impa.br